

SNLS: Overview and High-z Spectroscopy

D. Andrew Howell for the SNLS Collaboration

*Department of Astronomy and Astrophysics, The University of Toronto,
60 St. George Street, Toronto, ON, Canada*

Abstract. The Supernova Legacy Survey (SNLS)¹ will discover and obtain $g'r'i'z'$ lightcurves for more than 700 spectroscopically confirmed SNe Ia ($0.3 < z < 0.9$) to differentiate between competing models for Dark Energy. We fit the multicolor lightcurves of the candidates to determine which are likely SNe Ia and send them for follow-up spectroscopy to the Keck, VLT, and Gemini telescopes. Here we show the results from Gemini, where we send most of our highest redshift ($0.6 < z < 0.9$) targets.

1. Lightcurves

As a part of the Canada-France-Hawaii Telescope Legacy Survey (CFHTLS), the SNLS uses a rolling search strategy to discover and monitor SNe in four one degree fields over the course of five years. SNe are observed in $g'r'i'z'$, allowing unprecedented measurement of reddening and SN intrinsic colors. Observations are made every 2-3 rest frame days during dark and gray time. A typical run includes 10 observing blocks of 5 visits to each of two deep fields. Because we always observe the same fields, every visit brings new SN discoveries and simultaneously adds points to existing lightcurves. See the contribution of Sullivan et al. to these proceedings for an overview of the SNLS.

2. Spectroscopy

For the most promising candidates, spectroscopy on 8-10m telescopes must be obtained to determine the redshift and SN type. We prescreen candidates – early data are fit with a Ia lightcurve and poor fits (core-collapse SNe or AGN) are rejected. For good fits (likely SNe Ia), the redshift and phase are estimated, allowing optimal scheduling of observations and instrument setup.

Spectroscopy of candidate SNe is carried out on Gemini, Keck and VLT. The highest redshift targets are observed at Gemini N and S with GMOS (Hook et al. 2004), where we make use of the Nod and Shuffle mode to virtually eliminate the systematic errors associated with the subtraction of sky lines in the red. Gemini also observes some lower redshift targets in classical mode in the D3 (Extended Groth Strip) field, which cannot be seen by VLT. All GMOS observations use the R400 grating, $0.75''$ slit, and either the 680 or 720nm central wavelength setup. The CCD is binned 2×2 .

¹For a complete author list and candidates see <http://cfht.hawaii.edu/SNLS>

3. Identification

Spectra are fit with a χ^2 matching program (Howell & Wang 2002) against a library of SNe templates. Template host galaxy light is also fit and subtracted, and the redshift of the SN is estimated. This permits a determination of the SN type even when it is buried in a host galaxy. An expert must still make the final decision, but the quantitative comparison of hundreds of SN spectra offers robustness, accuracy, and speed advantages over unaided human classification.

4. Results

At Gemini, in semesters 2003B and 2004A we observed 48 targets, 29 of which are consistent with a SN Ia classification. We show some example spectra in Figure 1.

Acknowledgments. The SNLS collaboration acknowledges the CFHT Queue Service Observing team, the agencies CNRS/IN2P3, CNRS/INSU and CEA in France, and NSERC and CIAR in Canada. Based on observations obtained at the Gemini Observatory, and with Mega Prime, a joint project of CFHT, CEA/DAPNIA, and HIA.




Figure 1: Example Gemini spectra of confirmed Type Ia SNLS SNe from 2003B/04A. We show data after host galaxy subtraction (if necessary), re-binned to $5\text{\AA}$, with smoothed data overplotted. The best-fitting SN template is shown underneath.

References

- Hook, I. et al. 2004, PASP, 116, 425
 Howell, D. A. & Wang, L. 2002, AAS, 201, 9103